



# ANALECTA MEDICA



Atheroma  
of the  
coronary  
artery

Media  
Internal elastic  
membrane  
Intima enlarged  
by atheromatous  
growth



Reduced facsimiles

**DIGOXIN** (B. W. & Co.)

(See pages 10 and 11)

# ANALECTA MEDICA

SOME pharmaceutical products of special interest are described in the following pages. They have been selected as representative of the preparations which have gained for Burroughs Wellcome & Co. a world-wide reputation for the purity, accuracy and reliability of their products.

The preparations described reflect many important phases of modern medicine. The new chemotherapy is represented by Sulphonamide-P (*pages 24-25*), which, in spite of many recently-introduced elaborations of the simple sulphanilamide molecule, remains at once the most economical and the most generally useful compound of the sulphonamide group. Chemical synthesis has entered also into the field of hormone therapy, and, in the case of Stilboestrol (*pages 22-23*), has provided the clinician with a product which is not only at least as active as the natural oestrogens which it replaces, but possesses the advantage of being effective by oral administration as well as by injection.

In certain instances the space available in this booklet is not sufficient for more than a brief outline of the technique of administration and other details; in such cases the reader is referred to Wellcome's Medical Diary and to special literature which will gladly be forwarded on request. Attention is directed also to the comprehensive index on *pages 26-28* of this booklet, which facilitates reference to individual products.

Mo: pam - wets.

QV 26

193\*

B97a2



22500886770



| PRODUCT                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | INDICATIONS                                                                                                                                                                                                                     |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|-------------------------|-------|------------------------|-------|-----------------------------|-------|-----------------|-------|-----------------------|------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Adrenaline Tartrate</b><br/> <b>Solution of</b><br/> <small>TRADE MARK</small> <b>'Wellcome'</b> <small>BRAND</small><br/> (Adrenaline, 1 in 1000)</p>                                                                                                                                                                                                                                                                                                                                                  | <p>As a hæmostatic in epistaxis, local capillary hæmorrhage and hæmatemesis; also (applied as a spray) in acute coryza, hay fever and congestive conditions of the nasal mucous membrane</p>                                    |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| <p><b>Adrenaline</b><br/> 1 in 1000<br/> <small>TRADE MARK</small> <b>'Hypoloid'</b> <small>BRAND</small></p>                                                                                                                                                                                                                                                                                                                                                                                                 | <p>Anaphylactic shock and other serum reactions; vaso-motor disturbances following arsphenamine administration; hypoglycæmia following insulin over-dosage; acute cardiac failure; anæsthetic collapse; paroxysms of asthma</p> |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| <p><b>Adrenaline</b><br/> <b>Solution of</b> (1 in 100)<br/> <small>TRADE MARK</small> <b>'Vaporole'</b> <small>BRAND</small></p>                                                                                                                                                                                                                                                                                                                                                                             | <p>Asthma</p>                                                                                                                                                                                                                   |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| <p><small>TRADE MARK</small> <b>'Alepol'</b> <small>MARK</small><br/> The Sodium Salts of a Selected Fraction of the less irritating lower melting-point Fatty Acids of Hydnocarpus Oil</p>                                                                                                                                                                                                                                                                                                                   | <p>Leprosy<br/> Causes local reactions in the leprotic tissue, resulting in release of the bacilli and their subsequent destruction</p>                                                                                         |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| <p><b>Alkaline Compound</b><br/> <b>Effervescent</b><br/> <small>TRADE MARK</small> <b>'Tabloid'</b> <small>BRAND</small></p> <p>R</p> <table> <tr> <td>Sodii Bicarbonatis,</td> <td>gr. 5</td> </tr> <tr> <td>Calcii Lactophosphatis,</td> <td>gr. 3</td> </tr> <tr> <td>Potassii Bicarbonatis,</td> <td>gr. 1</td> </tr> <tr> <td>Magnesii Sulphatis, Anhyd.,</td> <td>gr. 1</td> </tr> <tr> <td>Sodii Chloridi,</td> <td>gr. 1</td> </tr> <tr> <td>Salis Effervescentis,</td> <td>q.s.</td> </tr> </table> | Sodii Bicarbonatis,                                                                                                                                                                                                             | gr. 5 | Calcii Lactophosphatis, | gr. 3 | Potassii Bicarbonatis, | gr. 1 | Magnesii Sulphatis, Anhyd., | gr. 1 | Sodii Chloridi, | gr. 1 | Salis Effervescentis, | q.s. | <p>As an alkaliniser in nausea and vomiting of pregnancy, cyclical vomiting in children, post-anæsthetic acidosis, acute infections, pyrexial conditions and other disorders involving acidosis; and as an alkaline diuretic in pyelitis, cystitis, urethritis and allied conditions</p> |
| Sodii Bicarbonatis,                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | gr. 5                                                                                                                                                                                                                           |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| Calcii Lactophosphatis,                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | gr. 3                                                                                                                                                                                                                           |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| Potassii Bicarbonatis,                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | gr. 1                                                                                                                                                                                                                           |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| Magnesii Sulphatis, Anhyd.,                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | gr. 1                                                                                                                                                                                                                           |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| Sodii Chloridi,                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | gr. 1                                                                                                                                                                                                                           |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| Salis Effervescentis,                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | q.s.                                                                                                                                                                                                                            |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |
| <p><b>Amyl Nitrite</b><br/> <small>TRADE MARK</small> <b>'Vaporole'</b> <small>BRAND</small><br/> Min. 3 and min. 5.</p>                                                                                                                                                                                                                                                                                                                                                                                      | <p>Paroxysms of angina pectoris; aneurysmal pain; asthma; anæsthetic and other states of cardiac or respiratory collapse</p>                                                                                                    |       |                         |       |                        |       |                             |       |                 |       |                       |      |                                                                                                                                                                                                                                                                                          |

## HOW TO USE

For local application, use in strengths of 1 in 100,000 to 1 in 1000; orally, 1/2 to 1 fluid ounce. Adrenaline tartrate is considerably more stable than the hydrochloride, and its solution is less acid

By intravenous injection, 1 c.c. in 500 c.c. (about a pint) of normal saline. By subcutaneous injection, 5 to 15 minims. The official dose is from 2 to 8 minims

To be inhaled by the mouth from an 'AGLA' or other suitable atomiser  
Not for injection

One to five c.c. of 3 per cent. solution, intramuscularly or subcutaneously, or one to five c.c. of 1 per cent. solution, intravenously. (*See special literature*)

One or two, dissolved in half-a-tumblerful of water, after meals. Each product represents, in alkalinising effect, about 21 grains of bicarbonate

The delicate glass capsule (surrounded by absorbent material and silken netting) should be crushed and the vapour inhaled

## PACKINGS

**'Wellcome' Solution of Adrenaline Tartrate**, amber-coloured, glass-stoppered bottles of 10 c.c. and 25 c.c.

**'Hypoloid' Adrenaline (1 in 1000)**, boxes of 10 'Hypoloid' ampoules of 1 c.c., and rubber-capped bottles of 10 c.c.

**'Vaporole' Solution of Adrenaline (1 in 100)**, amber-coloured, glass-stoppered bottles of 5 c.c.

**'Alepol,'** bottles of 25 gm. and 100 gm.

**'Tabloid' Alkaline Compound**, Effervescent, tubes of 25

**'Vaporole' Amyl Nitrite**, each strength in boxes of 12

| PRODUCT                                                                                                                                                                  | INDICATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Ascorbic Acid</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>0.005 gm. and 0.025 gm.</p> <p>Containing 100 and 500 International Units of Vitamin C respectively</p> | <p>For the prophylaxis and treatment of scurvy and other manifestations of Vitamin C deficiency</p>                                                                                                                                                                                                                                                                                                                                               |
| <p>TRADE 'Avenyl' MARK</p> <p>2-Myristoxy-mercuri-3-hydroxy-benzaldehyde</p> <p>A mercury preparation soluble in Hydnocarpus Oil and in 'Moogrol'</p>                    | <p>Leprosy complicated by syphilis. Controls the syphilis and renders W.R. negative without causing reaction in the leprous element of the condition</p>                                                                                                                                                                                                                                                                                          |
| <p><b>Barbitone</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>Gr. 5 and 0.5 gm.*</p>                                                                                      | <p>Nervous insomnia; delirium of fevers; excitable mental states; delirium tremens; migraine; sea-sickness</p>                                                                                                                                                                                                                                                                                                                                    |
| <p><b>Barbitone, Soluble</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>Gr. 5</p>                                                                                          | <p>As for 'Tabloid' Barbitone (<i>above</i>). More rapid in action, owing to greater solubility</p>                                                                                                                                                                                                                                                                                                                                               |
| <p>TRADE MARK 'Bicreol' BRAND</p> <p><b>Bismuth Cream</b></p> <p>0.15 gm. metallic bismuth in 1 c.c., in a creo-camph base</p>                                           | <p><b>Syphilis</b></p> <p>Bismuth and its salts, injected intramuscularly, exert a powerful spirillicidal action, only slightly less effective than the arsphenamines, and superior to mercury. Bismuth preparations may be given in courses of one injection weekly for eight to twelve weeks, until a minimum of 2 grammes of bismuth has been given, the courses alternating with courses of arsphenamine over a period of about two years</p> |
| <p><b>Bismuth Metal</b></p> <p>TRADE MARK 'Hypoloid' BRAND</p> <p>0.2 gm. in 1 c.c., suspended in a sterile isotonic glucose solution</p>                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <p><b>Bismuth Oxychloride</b></p> <p>TRADE MARK 'Hypoloid' BRAND</p> <p>0.1 gm. in 1 c.c., suspended in a sterile isotonic saline solution</p>                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                   |

\* Obtainable in the British Empire Overseas to special order only

## HOW TO USE

## PACKINGS

Prophylactic dose: 0.025 gramme to 0.05 gramme (500 to 1000 units) daily. Therapeutic dose: 0.1 gramme to 0.25 gramme (2000 to 5000 units) daily

**'Tabloid' Ascorbic Acid**, each strength in bottles of 25

A 0.25 per cent. solution in Hydnocarpus Oil, or a 0.5 per cent. saturated solution in 'Moogrol,' is injected subcutaneously, commencing with 1 c.c., increased to 4 c.c. or more. (*See special literature*)

**'Avenyl,'** tubes of 1 gm.

Gr. 5 to gr. 10. The maximum official dose is gr. 10; full doses should not be given until the patient's susceptibility to smaller doses has been determined

**'Tabloid' Barbitone**, gr. 5 in bottles of 25 and 100; 0.5 gm. in bottles of 25

As for **'Tabloid' Barbitone** (*above*)

**'Tabloid' Barbitone, Soluble**, bottles of 25 and 100

By intramuscular injection. The suggested commencing dose is 1 c.c., increased to 2 c.c.

**'Bicreol,'** glass jars containing 1/2 oz. and 2 oz.

By intramuscular injection. The suggested commencing dose is 1 c.c., increased to 1.5 c.c.

**'Hypoloid' Bismuth Metal**, rubber-capped bottles of 5 c.c., 10 c.c. and 25 c.c.

By intramuscular injection. The suggested commencing dose is 1 c.c., increased to 1.5 c.c.

**'Hypoloid' Bismuth Oxychloride**, rubber-capped bottles of 25 c.c.

| PRODUCT                                                                                                                                                                                          | INDICATIONS                                                                                                                                                                                                         |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Blaud Pill</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>Gr. 5, gr. 10 and 0.25 gm.*</p> <p>(Sugar-coated)</p> <p>Each represents not less than 22.5 per cent. of ferrous carbonate</p>     | <p>Nutritional and secondary anæmias; amenorrhœa; convalescence. 'Tabloid' Blaud Pill combines a high degree of effectiveness as a hæmatinic with almost complete absence of gastric irritation or constipation</p> |
| <p><b>Calciferol</b></p> <p>The pure crystalline anti-rachitic principle, with a constant activity of 40,000 units per mgm.</p>                                                                  | <p>For the prophylaxis and treatment of rickets (<i>see also</i> 'Tabloid' Calcium Gluconate with Calciferol, <i>below</i>)</p>                                                                                     |
| <p><b>Calcium Gluconate</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>Gr. 20</p> <p>(Chocolate-flavoured Base)</p>                                                                                | <p>During pregnancy, to furnish dietary calcium; urticaria, dermatitis herpetiformis and other skin conditions; hay fever; chilblains: and as a pre-operative measure</p>                                           |
| <p><b>Calcium Gluconate with Calciferol</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>R. Calci Gluconatis, 0.325 gm.<br/>Calciferolis, 0.000025 gm.</p>                                           | <p>For administration during pregnancy and lactation, as a prophylactic against rickets in the growing foetus or infant</p>                                                                                         |
| <p><b>Calcium Gluconate and Iron (Effervescent)</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>R. Calci Gluconatis, gr. 20<br/>Ferri et Ammonii Citratis, gr. 3<br/>Salis Effervescentis, q.s.</p> | <p>For administration during pregnancy and lactation, and at other times, as a prophylactic against calcium and iron deficiency</p>                                                                                 |
| <p><b>Calcium Lævulate</b></p> <p>TRADE MARK 'Hypoloid' BRAND</p> <p>Ampoules of 1 gm. in 10 c.c.</p>                                                                                            | <p>Tetany; vomiting of intestinal obstruction; coeliac rickets; as a pre-operative measure; and all conditions where parenteral calcium therapy is indicated</p>                                                    |

\* Obtainable in the British Empire Overseas to special order only



## HOW TO USE

## PACKINGS

One, of the required strength, gradually increased if necessary, taken with a little water, twice or thrice daily, after food. The maximum official dose is thirty grains

**'Tabloid' Bland Pill**, each strength in bottles of 100. (Numerous combinations of **'Tabloid' Bland Pill** with various hæmatinic and alterative medicaments are also available; *see* Wellcome's Medical Diary)

Prophylactic dose (for infants): 0·000025 gm. to 0·00005 gm. (1000 to 2000 units) daily.  
Therapeutic dose (for infants): 0·00005 gm. to 0·000075 gm. (2000 to 3000 units) daily

**'Tabloid' Calciferol**, 0·0001 gm. (4000 units) and 0·00025 gm. (10,000 units), each strength in bottles of 25 and 100. **'Wellcome' Solution of Calciferol** (3000 units in 1 gm.), bottles of 1 fl. oz. (*with dropper*)

Adults: two to three products, three times a day, before food. Children: one product, three times a day, before food. The maximum official dose is sixty grains

**'Tabloid' Calcium Gluconate**, tubes of 25

One, or more, to be taken with water, daily. Each product contains 1000 International Units of Vitamin D

**'Tabloid' Calcium Gluconate with Calciferol**, bottles of 25 and 100

One or two products, daily, in a large tumblerful of water

**'Tabloid' Calcium Gluconate and Iron (Effervescent)**, tubes of 20

By intravenous injection. Adults: 10 c.c. daily, or every second day. Children: 2 c.c. to 5 c.c., according to age

**'Hypoloid' Calcium Lævulate**, boxes of 5 **'Hypoloid'** ampoules

| PRODUCT                                                                                                                                                                                                                                                                                                      | INDICATIONS                                                                                                                                                                                   |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Calcium Mandelate Compound</b></p> <p>TRADE MARK 'Wellcome' BRAND</p> <p>A pleasantly-flavoured powder, readily suspensible in water or milk.</p>                                                                                                                                                      | <p>Urinary tract infections, including <i>Streptococcus faecalis</i></p>                                                                                                                      |
| <p><b>Carbarson</b></p> <p><i>p</i>-Carbamidophenylarsonic Acid</p> <p>A stable arsenical compound containing 28.85 per cent. of arsenic</p>                                                                                                                                                                 | <p>Chronic intestinal amoebiasis. Of low toxicity; does not produce nausea or vomiting</p>                                                                                                    |
| <p><b>Carotene</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>0.002 gm. (Sugar-coated)</p> <p>A mixture of <math>\alpha</math>-, <math>\beta</math>- and <math>\gamma</math>-carotene, obtained from various vegetable sources</p>                                                                             | <p>Xerophthalmia and night blindness; keratinisation of upper respiratory epithelium; cutaneous lesions and keratosis; and other conditions arising from Vitamin A deficiency</p>             |
| <p>TRADE MARK 'Diginutin' MARK</p> <p>A stable solution of the total glucosides of <i>Digitalis purpurea</i> leaf (physiologically standardised) 1 International Unit in 1 c.c.</p> <p>Also issued as 'Tabloid' 'Diginutin,' min. 5 and min. 10 (containing 0.3 and 0.6 International Unit respectively)</p> | <p>Auricular fibrillation or flutter; congestive heart failure; paroxysmal nocturnal dyspnoea; mild condition of senile heart; and circulatory disturbances associated with renal disease</p> |
| <p><b>Digitalis Leaf, B.P.</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>Gr. 1/2 and gr. 1 (containing 0.3 and 0.6 International Unit respectively)</p>                                                                                                                                                       | <p>As for 'Diginutin' (<i>above</i>)</p>                                                                                                                                                      |

## HOW TO USE

## PACKINGS

A level measureful (4.4 grammes, equivalent to 3 grammes of Mandelic Acid), to be taken in four tablespoonfuls of water or milk, four times a day. (*See special literature*)

**'Wellcome' Calcium Mandelate Compound**, amber-coloured bottles of 125 grammes. A measure and a supply of test papers are provided with each packing

**'Tabloid' Carbarsone**: one, by mouth, twice daily, for ten days; repeated, if desired after a rest period of a week to ten days. **'Wellcome' Carbarsone**: two grammes, dissolved in 200 c.c. of 2 per cent sodium bicarbonate solution, administered per rectum on alternate nights for a maximum of five doses

**'Tabloid' Carbarsone**, 0.25 gm., bottles of 20 and 100†; ‡**'Wellcome' Carbarsone**, bottles of 25 grammes and 100 grammes

One to five products, according to age, three times a day. Each product is equivalent to 2000 International Units of Vitamin A

**'Tabloid' Carotene**, bottles of 25

For oral administration. 25 minims every six hours, and effect carefully watched for in five to seven days. Maintenance dose, five to fifteen minims

**'Diginutin'**, bottles of 1 fl. oz. and 8 fl. oz.; **'Tabloid' 'Diginutin'**, each strength in bottles of 25 and 100

One to three of the gr. 1/2 strength every six hours, and effect carefully watched for in five to seven days, or a single dose of gr. 3 to gr. 10

**'Tabloid' Digitalis Leaf**, each strength in bottles of 25 and 100

† Supplied in the British Isles, India and China only

‡ Supplied in the British Isles and China only

| PRODUCT                                                                                                                                                                                                                                                                                                      | INDICATIONS                                                                                                                                                                               |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Digoxin (B. W. &amp; Co.)</b><br/>(Under British Patent)</p> <p>A pure, stable, crystallised glucoside, isolated from the leaves of <i>Digitalis lanata</i></p> <p>'Tabloid' Digoxin, 0.25 mgm.; Solution of Digoxin (B. W. &amp; Co.), 0.5 mgm. in 1 c.c.; 'Hypoloid' Digoxin, 0.5 mgm. in 1 c.c.</p> | <p>May be used whenever digitalis therapy is indicated. Oral administration produces rapid and consistent response; intravenous injection necessary only in exceptional circumstances</p> |
| <p>TRADE 'Emocin' MARK</p> <p><b>Throat Lozenge</b></p> <p>TRADE 'Tabloid' BRAND</p> <p>Each product contains 2 grains 'Empirin' Acetylsalicylic Acid, combined with a flavoured demulcent base</p>                                                                                                          | <p>For use after tonsillectomy, and in inflammatory and irritative conditions of the throat</p>                                                                                           |
| <p>TRADE 'Empirin' MARK</p> <p>(ACETYSALICYLIC ACID)</p> <p><b>Compound</b></p> <p>TRADE 'Tabloid' BRAND</p> <p>R 'Empirin' Acid.<br/>Acetylsalicyl., gr. 3-1/2<br/>Phenacetini, gr. 2-1/2<br/>Caffeinæ, gr. 1/2</p>                                                                                         | <p>As an analgesic in neuralgia, migraine, toothache, rheumatism, sciatica, dysmenorrhœa, lumbago, etc., and as an antipyretic in coryza, influenza and other febrile states</p>          |
| <p>TRADE 'Empirin' MARK</p> <p>(ACETYSALICYLIC ACID)</p> <p><b>Compound with Codeine</b></p> <p>TRADE 'Tabloid' BRAND</p> <p>R 'Empirin' Acid.<br/>Acetylsalicyl., gr. 3-1/2<br/>Phenacetini, gr. 2-1/2<br/>Caffeinæ, gr. 1/2<br/>Codeinæ Phosphatis, gr. 1/8</p>                                            | <p>As for 'Tabloid' 'Empirin' Compound (<i>above</i>). The inclusion of codeine phosphate considerably increases the analgesic action of the product</p>                                  |



## HOW TO USE

## PACKINGS

Orally, 0.25 mgm. to 1.5 mgm.  
By intravenous injection,  
0.5 mgm. to 1 mgm. Care must  
be taken to avoid extravenous  
injection, as the solution is  
irritant

**'Tabloid' Digoxin**, bottles of  
25 and 100. **Solution of  
Digoxin (B. W. & Co.)**, bottles  
of 10 c.c. and 30 c.c. (supplied  
with pipette) and 250 c.c.  
**'Hypoloid' Digoxin**, boxes of  
10 ampoules

One, slowly dissolved in the  
mouth and repeated as required.  
Following tonsillectomy, one  
product may be given about  
ten minutes before each meal  
for three or four days

**'Tabloid' 'Emocin,'** tubes of 25

One to two, taken with a little  
water

**'Tabloid' 'Empirin' Compound,**  
bottles of 25 and 100

One to two, taken with a little  
water

**'Tabloid' 'Empirin' Compound  
with Codeine**, bottles of 25  
and 100

| PRODUCT                                                                                                                                                                                                                                                                                                                                                                                                             | INDICATIONS                                                                                                                                                                                                                                                                   |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Ephedrine Hydrochloride</b><br/> <small>TRADE MARK</small> <b>'Tabloid'</b> <small>BRAND</small></p> <p>Gr. 1/4, gr. 1/2 and 0.03 gm.*</p>                                                                                                                                                                                                                                                                    | <p>Asthma, hay fever, urticaria and allergic conditions generally; narcolepsy; enuresis; myasthenia gravis; post-encephalitic Parkinsonism</p>                                                                                                                                |
| <p><b>Ephedrine Hydrochloride</b><br/> <small>TRADE MARK</small> <b>'Hypoloid'</b> <small>BRAND</small></p> <p>0.03 gm. in 1 c.c.</p>                                                                                                                                                                                                                                                                               | <p>Asthma, circulatory failure, and other urgent indications</p>                                                                                                                                                                                                              |
| <p><b>Ephedrine Compound</b><br/> <small>TRADE MARK</small> <b>'Elixoid'</b> <small>BRAND</small></p> <p>A pleasantly-flavoured preparation, containing Ephedrine Hydrochloride, gr. 1/4; Syrup of Tolu, min. 15; Tincture of Virginian Prune, min. 2; Chloroform, min. 1/8, in each fluid drachm</p>                                                                                                               | <p>Whooping cough. In severe cases, the effect may be enhanced by the concurrent administration of belladonna</p>                                                                                                                                                             |
| <p><b>Ephedrine Spray Compound</b><br/> <small>TRADE MARK</small> <b>'Vaporole'</b> <small>BRAND</small></p> <p>Contains Ephedrine, 0.9 per cent.; Menthol, Camphor and Oil of Thyme, of each 0.5 per cent., in a base of 'Paroleine' Liquid Paraffin</p>                                                                                                                                                           | <p>For local application in hay fever and other engorged and catarrhal conditions of the naso-pharynx</p>                                                                                                                                                                     |
| <p><b>Ergometrine</b></p> <p>A crystalline alkaloid isolated from aqueous extract of ergot</p> <p>Oral: 'Tabloid' Ergometrine, 0.5 mgm.; 'Wellcome' Solution of Ergometrine, 0.5 mgm. in 2.5 c.c. Intramuscular: 'Tabloid' Hypodermic Ergometrine, 0.5 mgm.; 'Hypoloid' Ergometrine, 0.5 mgm. in 1 c.c. Intravenous: 'Tabloid' Hypodermic Ergometrine, 0.125 mgm.; 'Hypoloid' Ergometrine, 0.125 mgm. in 1 c.c.</p> | <p>For administration after parturition and during the puerperium. Suggested as a useful means of treating post-partum hæmorrhage, particularly in cases where pituitary extract may produce alarming pallor and shock. May be of value in migraine (given intravenously)</p> |

\* Obtainable in the British Empire Overseas to special order only

## HOW TO USE

## PACKINGS

For oral or hypodermic administration. The maximum official dose is one-and-a-half grains

**'Tabloid' Ephedrine Hydrochloride**, each strength in bottles of 25 and 100; gr. 1/2 also in tubes of 6

One to three c.c., injected hypodermically

**'Hypoloid' Ephedrine Hydrochloride**, boxes of 10 'Hypoloid' ampoules

One to two teaspoonfuls, or more, as required

**'Elixoid' Ephedrine Compound**, bottles containing 4 fl. oz. and 16 fl. oz.

To be used with a 'Paroleine' or other suitable atomiser, or a few drops instilled into the nostrils by means of the dropper supplied with the smaller size

**'Vaporole' Ephedrine Spray Compound**, bottles of 1 fl. oz. (supplied with dropper) and 16 fl. oz.†

Orally, 0.5 mgm. to 1 mgm.  
Intramuscularly, 0.5 mgm., repeated in one hour if desired.  
Intravenously, 0.125 mgm., repeated in one hour if desired

**'Tabloid' Ergometrine**, bottles of 25 and 100. **'Wellcome' Solution of Ergometrine**, bottles of 30 c.c. (with pipette). **'Tabloid' Hypodermic Ergometrine**, each strength in tubes of 12. **'Hypoloid' Ergometrine**, each strength in boxes of 10 ampoules

† Not supplied in China or India

| PRODUCT                                                                                                                                                                                                                                                                             | INDICATIONS                                                                                                                                                                      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Ergotoxine Ethanesulphonate</b></p> <p>'Tabloid' Ergotoxine Ethanesulphonate, 0.0005 gm.; 'Tabloid' Hypodermic Ergotoxine Ethanesulphonate, gr. 1/100; 0.00025 gm.; and 0.0005 gm.; 'Wellcome' Ergotoxine Ethanesulphonate, B.P.</p>                                          | <p>As a routine after labour, to assist involution of the uterus and to reduce the risk of post-partum hæmorrhage; and in menorrhagia and other types of uterine hæmorrhage</p>  |
| <p>TRADE 'Ernutin' MARK</p> <p>A clear solution, presenting Ergotoxine—together with 'Ergamine' and 'Tyramine'—in definite amount and in a pure state</p> <p>'Ernutin' (Oral) contains 0.033 per cent., and 'Hypoloid' 'Ernutin' 0.1 per cent., of Ergotoxine Ethanesulphonate.</p> | <p>As for Ergotoxine Ethanesulphonate (<i>above</i>)</p>                                                                                                                         |
| <p><b>Erythrityl Tetranitrate</b></p> <p>TRADE 'Tabloid' BRAND</p> <p>Gr. 1/4, gr. 1/2 and gr. 1</p>                                                                                                                                                                                | <p>Angina pectoris; arterial hypertension; and other conditions where vaso-dilator action is required. Effect maximal in about one hour, and lasts approximately three hours</p> |
| <p>TRADE 'Ethidol' BRAND</p> <p><b>Ethyl Iodo-ricinoleate</b></p> <p>Non-irritating; non-staining. Contains 20 per cent. of Iodine in organic combination</p>                                                                                                                       | <p>Subcutaneous inflammation; joint or muscle traumata; enlarged cervical glands; orchitis; lumbago and other myalgias</p>                                                       |
| <p>TRADE 'Eulykol' MARK</p> <p>Phenylethyl Esters of a Selected Fraction of the Acids of Hydno-carpus Oil—sometimes designated "Phenylethyl Hydno-carpate"</p>                                                                                                                      | <p>Lupus vulgaris; leprosy</p>                                                                                                                                                   |
| <p><b>Ferrous Chloride Citrated</b></p> <p>TRADE 'Tabloid' BRAND</p> <p>Gr. 3 (Sugar-coated)</p>                                                                                                                                                                                    | <p>Nutritional and secondary anæmias; amenorrhœa; convalescence</p>                                                                                                              |



## HOW TO USE

## PACKINGS

By intramuscular or subcutaneous injection, 0.0005 gm. to 0.001 gm. (gr. 1/120 to gr. 1/60). Also given orally

'Tabloid' Ergotoxine Ethanesulphonate, bottles of 25.  
'Tabloid' Hypodermic Ergotoxine Ethanesulphonate, each strength in tubes of 12.  
'Wellcome' Ergotoxine Ethanesulphonate, B.P., tubes of 0.1 gm.

'Ernutin' (Oral), 30 to 60 minims. 'Hypoloid' 'Ernutin,' by subcutaneous or intramuscular injection, 0.3 c.c. to 0.6 c.c.

'Ernutin' (Oral), bottles of 30 c.c., 4 fl. oz. and 16 fl. oz.  
'Hypoloid' 'Ernutin,' 0.6 c.c., boxes of 6 ampoules

One, or more, of the gr. 1/4 or gr. 1/2 strength, repeated as may be necessary. The maximum official dose is one grain, but the smallest dose should always be used at first

'Tabloid' Erythrityl Tetranitrate, gr. 1/4 and gr. 1/2, bottles of 25 and 100; gr. 1, bottles of 12

Inunction: Rub lightly into the part until absorbed. Intraglandular injection: Half c.c. to one c.c. on alternate days

'Ethidol,' bottles of 1 fl. oz. and 4 fl. oz.

Initial dose: 0.1 c.c., injected intradermally. One week later the whole patch may be infiltrated with 10 doses of 0.1 c.c. at separate spots. This dose may be repeated weekly

'Eulykol,' bottles of 25 c.c.

One to two, taken with copious draughts of water, after food. The maximum official dose is five grains

'Tabloid' Ferrous Chloride, Citrated, bottles of 100

| PRODUCT                                                                                                                                                                                                                                    | INDICATIONS                                                                                                                                                                                                                         |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Glyceryl Trinitrate</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>(Trinitrin)</p> <p>Gr. 1/200; gr. 1/100; gr. 1/50;<br/>and 0.0005 gm.*</p>                                                                                          | <p>Used to relieve attacks of angina pectoris, of weak and irregular heart action, and of asthma. May be effective in migraine (preferably with strychnine)</p>                                                                     |
| <p>TRADE MARK 'Infundin' BRAND</p> <p><b>Pituitary</b><br/>(Posterior Lobe)<br/>Extract</p> <p>10 International Units per c.c. (identical with Liquid Extract of Pituitary, B.P. 1932), and 5 International Units per c.c.</p>             | <p>Post-partum hæmorrhage; uterine inertia (should not be given before the os is about one-half dilated); intestinal paresis (to initiate peristalsis); and diabetes insipidus</p>                                                  |
| <p><b>Insulin</b></p> <p>TRADE MARK 'Wellcome' BRAND</p> <p>Made with pure crystalline insulin</p> <p>Also issued: 'Tabloid' Hypodermic Insulin Hydrochloride (<i>Sterile</i>). Each product presents a dose of 10 International Units</p> | <p>Diabetes mellitus; malnutrition</p>                                                                                                                                                                                              |
| <p><b>Protamine Insulin</b><br/>(with Zinc)—Suspension</p> <p>TRADE MARK 'Wellcome' BRAND</p> <p>40 Units and 80 Units per c.c.</p>                                                                                                        | <p>Retarded action renders it specially suitable for severe and complicated cases of diabetes; may also be used to reduce the number of injections in less severe cases</p>                                                         |
| <p>TRADE MARK 'Iodicin' MARK</p> <p><b>Calcium Iodo-ricinoleate</b></p> <p>Contains 33-1/3 per cent. of Iodine in organic combination</p> <p>'Tabloid' 'Iodicin,' gr. 3 (<i>Capsule</i>), and 0.03 gm. (<i>Chocolate Base</i>)</p>         | <p>In all conditions where iodide therapy is indicated; may be given over long periods without causing symptoms of iodism. The 0.03 gm. (<i>Chocolate Base</i>) product is particularly suitable for administration to children</p> |

\* Obtainable in the British Empire Overseas to special order only

## HOW TO USE

## PACKINGS

One of the gr. 1/200, gr. 1/100, or 0.0005 gm. strength, occasionally, as necessity arises. During anginal attacks, gr. 1/50, chewed or dissolved in the mouth. In general, gr. 1/200 every hour, gr. 1/100 every two hours, or gr. 1/50 every three hours

**'Tabloid' Glyceryl Trinitrate**, each strength in bottles of 25 and 100

Half to two c.c. of the required strength, injected intramuscularly, or intravenously in normal saline. Obstetrical: Ante-partum, 2 units; post-partum, 10 to 15 units, in both cases by intramuscular injection. B.P. dose, two to five units by subcutaneous injection

**'Hypoloid' 'Infundin,'** 10 International Units per c.c., ampoules of 0.5 c.c. and 1 c.c., each in boxes of 6 and 25. 5 International Units per c.c., ampoules of 0.5 c.c. and 1 c.c., each in boxes of 6

Five units, or more, by hypodermic injection, up to daily average of 10 to 30 units. Much larger doses are required in coma. The B.P. dose is 5 to 100 units

**'Wellcome' Insulin**, rubber-capped, amber-glass phials of 100 International Units in 5 c.c.; 200 International Units in 5 c.c.; 400 International Units in 5 c.c.\*; and 200 International Units in 10 c.c. **'Tabloid' Hypodermic Insulin Hydrochloride**, cartons containing 10 products in one tube

Usual single daily dose, 20 to 60 units. May be given with unmodified insulin (*see special literature*)

**'Wellcome' Protamine Insulin (with Zinc) Suspension**, rubber-capped, clear-glass phials of 200 International Units in 5 c.c.; 400 International Units in 5 c.c.; and 400 International Units in 10 c.c.

Orally, gr. 1/2 to gr. 9

**'Tabloid' 'Iodacin,'** gr. 3 (Capsule), boxes of 50; 0.03 gm. (Chocolate Base), bottles of 25 and 100

| PRODUCT                                                                                                                                                                      | INDICATIONS                                                                                                                                                                                                                                                                                                                                                                 |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>TRADE MARK <b>'Kepler'</b> MARK</p> <p><b>Cod Liver Oil</b><br/>with Malt Extract</p> <p>Presents essential vitamins in their natural and most effective state</p>        | <p>As a dietary adjunct in convalescence, during pregnancy and lactation, and for growing children. For list of combinations of 'Kepler' Cod Liver Oil with Malt Extract with various medicaments, <i>see</i> Wellcome's Medical Diary</p>                                                                                                                                  |
| <p>TRADE MARK <b>'Kharophen'</b> BRAND</p> <p><b>Acetarsol</b></p> <p>A pentavalent arsenical compound containing about 27·3 per cent. of arsenic in organic combination</p> | <p>Chronic amœbic dysentery (especially emetine-resistant forms); yaws; lamblia-sis; Vincent's angina; <i>Plasmodium vivax</i> infection</p>                                                                                                                                                                                                                                |
| <p><b>Mersalyl</b></p> <p>TRADE MARK <b>'Hypoloid'</b> BRAND</p> <p>A 10 per cent. sterile solution with 5 per cent. Theophylline</p>                                        | <p>Cardiac and renal œdema</p>                                                                                                                                                                                                                                                                                                                                              |
| <p>TRADE MARK <b>'Moogrol'</b> BRAND</p> <p><b>Mixture of Esters of Acids of the Chaulmoogric Series</b></p>                                                                 | <p>Leprosy</p>                                                                                                                                                                                                                                                                                                                                                              |
| <p><b>Iodised</b></p> <p>TRADE MARK <b>'Moogrol'</b> MARK</p> <p>A mixture of Esters of Acids of the Chaulmoogric Series with 0·5 per cent. of Iodine</p>                    | <p>Leprosy. The presence of iodine materially reduces the irritating tendency of the esters</p>                                                                                                                                                                                                                                                                             |
| <p>TRADE MARK <b>'Neokharsivan'</b> BRAND</p> <p><b>Neoarsphenamine</b></p>                                                                                                  | <p>Syphilis; yaws; anthrax 'Neokharsivan' and 'Khar-sulphan' are manufactured in England by Burroughs Wellcome &amp; Co., and are approved by the Ministry of Health for use under the Public Health (Venereal Diseases) Regulations. They comply with the requirements of the Therapeutic Substances Act (Great Britain), 1925 (1931 Regulations) and of the B.P. 1932</p> |
| <p>TRADE MARK <b>'Kharsulphan'</b> BRAND</p> <p><b>Sulpharsphenamine</b></p>                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                             |



## HOW TO USE

## PACKINGS

One teaspoonful to two dessert-spoonfuls, twice or thrice daily, after food

'Kepler' Cod Liver Oil with Malt Extract, bottles of two sizes

Gr. 4, twice a day (night and morning), after meals, for 10 days. This dosage should not be exceeded unless an interval of 7 to 14 days divides the courses

'Tabloid' 'Kharophen,' gr. 4, bottles of 25

By intravenous or intramuscular injection, 1/2 c.c. to 2 c.c.

'Hypoloid' Mersalyl, boxes of 10 ampoules of 1 c.c. and 5 ampoules of 2 c.c.

By intramuscular injection

'Moogrol,' bottles of 25 c.c., 100 c.c. and 1 litre‡

By intradermal and intramuscular injection, at weekly intervals (*for details of administration, see special literature*)

Iodised 'Moogrol,' bottles of 25 c.c., 100 c.c. and 1 litre‡

By intravenous injection in about 20 per cent. solution. The official dose of Neoarsphenamine is 0.15 gm. to 0.9 gm.

'Neokharsivan,' hermetically-sealed phials of 0.15 gm., 0.3 gm., 0.45 gm., 0.6 gm., 0.75 gm. and 0.9 gm.

By intramuscular or subcutaneous injection. The official dose of Sulpharsphenamine is 0.1 gm. to 0.6 gm.

'Kharsulphan,' hermetically-sealed phials of 0.15 gm., 0.3 gm., 0.45 gm. and 0.6 gm.

| PRODUCT                                                                                                                                                     | INDICATIONS                                                                                                                                                                                                      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>TRADE MARK 'Neostam' BRAND</p> <p><b>Stibamine Glucoside</b></p> <p>(The nitrogen-glucoside of Sodium <i>p</i>-Aminophenylstibonate)</p>                 | <p>Kala azar</p> <p>May also be of value in lympho-granuloma inguinale</p>                                                                                                                                       |
| <p>TRADE MARK 'Nicamide' BRAND</p> <p><b>Nicotinic Acid Diethylamide</b></p> <p>25 per cent. solution</p>                                                   | <p>As a respiratory and circulatory stimulant, in conditions associated with shock and depressed circulatory and respiratory states; asphyxia neonatorum; barbiturate poisoning; mental symptoms of pellagra</p> |
| <p><b>Nicotinic Acid</b></p> <p>Appears to be closely associated with, or derived from, the pellagra-preventing factor of Vitamin B<sub>2</sub></p>         | <p>Pellagra</p>                                                                                                                                                                                                  |
| <p><b>Phenobarbitone</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>Gr. 1/4, gr. 1/2, gr. 1 and gr. 1-1/2</p>                                                 | <p>Epilepsy; aural vertigo; insomnia of nervous origin</p>                                                                                                                                                       |
| <p><b>Phenobarbitone Soluble</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>Gr. 1/4, gr. 1/2, gr. 1 and gr. 1-1/2</p>                                         | <p>As for 'Tabloid' Phenobarbitone (<i>above</i>). More rapid in action, owing to greater solubility</p>                                                                                                         |
| <p>TRADE MARK 'Quinoxyl' BRAND</p> <p><b>Chiniofon</b></p> <p>Iodoxyhydroxyquinolinesulphonic acid, 80 per cent., with sodium bicarbonate, 20 per cent.</p> | <p>Amœbic dysentery</p>                                                                                                                                                                                          |

## HOW TO USE

## PACKINGS

0.1 gm. per 100 lb. body-weight, intravenously, as a 4 per cent. solution, on alternate days until 3 gm. per 100 lb. body-weight have been given (*see special literature*)

'Neostam,' hermetically-sealed phials of 0.05 gm., 0.1 gm., 0.2 gm., 0.3 gm., 0.4 gm., 0.5 gm. and 1 gm.

Orally, 1 c.c. to 2 c.c. By intravenous or intramuscular injection, 1 c.c. to 5 c.c.

'Nicamide' Nicotinic Acid Diethylamide, 25 per cent. solution, bottles of 15 c.c. (with pipette) and 100 c.c. 'Hypoloid' 'Nicamide,' 2 c.c. and 5 c.c., each in boxes of 6 ampoules

Orally, 0.1 gramme, 5 times daily. By injection, 0.05 gm. to 0.1 gm.

'Tabloid' Nicotinic Acid, 0.05 gm., bottles of 100. 'Hypoloid' Nicotinic Acid, 0.05 gm. in 1 c.c., boxes of 6 ampoules

Gr. 1/2 to gr. 2, orally. The maximum official dose is two grains

'Tabloid' Phenobarbitone, each strength in bottles of 25 and 100

As for 'Tabloid' Phenobarbitone (*above*)

'Tabloid' Phenobarbitone, Soluble, each strength in bottles of 25 and 100

Orally, 0.5 gm., three times a day (the official doses are 0.06 gm. to 0.5 gm.). By rectal injection, 200 c.c. of 2-1/2 per cent. solution, once daily, and retained (the official doses are 1 gm. to 5 gm.). 'Quinoxyl' is given by both routes in courses of 10 days, repeated if necessary in 5 to 7 days (*see special literature*)

'Quinoxyl,' bottles of 10 gm. and 100 gm. 'Tabloid' 'Quinoxyl,' 0.25 gm. and 0.5 gm., each strength in bottles of 25 and 100. 'Soloid' 'Quinoxyl,' 5 gm., tubes of 6

| PRODUCT                                                                                                                                                                                                                                                                                                      | INDICATIONS                                                                                                                                                                                                                                      |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Reduced Iron</b><br/> <small>TRADE MARK</small> <b>'Tabloid'</b> <small>BRAND</small><br/>           Gr. 2</p>                                                                                                                                                                                         | <p>Nutritional and secondary anæmias; amenorrhœa; convalescence</p>                                                                                                                                                                              |
| <p><small>TRADE MARK</small> <b>'Ryzamin-B'</b> <small>BRAND</small><br/> <b>Rice Polishings Concentrate</b><br/>           A clear, golden-brown syrupy preparation, with a potency of not less than 80 International Units of Vitamin B<sub>1</sub> per gramme</p>                                         | <p>Conditions associated with Vitamin B<sub>1</sub> deficiency, and as a dietary supplement for patients of all ages, to stimulate appetite and promote utilisation of food</p>                                                                  |
| <p><b>Sodium Nitrite Compound</b><br/> <small>TRADE MARK</small> <b>'Tabloid'</b> <small>BRAND</small><br/>           R<br/>           Sodii Nitritis, gr. 1/2<br/>           Erythritylis Tetranitratis, gr. 1/8<br/>           Mannitylis Hexanitratis, gr. 1/4<br/>           Sodii Hippuratis, gr. 1</p> | <p>Anginal pain; acute pulmonary œdema; angiospasm; tinnitus aurium; head throbbing; and other conditions where a vasodilator action is required. The product combines depressants whose promptness and duration of effect are complementary</p> |
| <p><b>Stilbœstrol</b><br/> <b>(Diethylstilbœstrol)</b><br/>           A synthetic œstrogen, closely resembling the natural hormone in therapeutic effect</p>                                                                                                                                                 | <p>Menopausal conditions, including atrophic vaginitis and kraurosis vulvæ; amenorrhœa; dysmenorrhœa; pruritus vulvæ</p>                                                                                                                         |
| <p><small>TRADE MARK</small> <b>'Stipolac'</b> <small>BRAND</small><br/> <b>Tetraiodophenolphthalein</b><br/> <b>'Stipolac'</b> Sodium Tetraiodophenolphthalein (<i>Iodophthalein</i>, <i>B.P.</i>) and <b>'Stipolac'</b> Acid Mixture, of each 3.5 grammes</p>                                              | <p>For radiographic examination of the gall-bladder</p>                                                                                                                                                                                          |
| <p><small>TRADE MARK</small> <b>'Stypven'</b> <small>BRAND</small><br/> <b>Russell Viper Venom</b><br/> <i>(Not for Injection)</i><br/>           Issued in rubber-stoppered bottles, accompanied by hermetically-sealed ampoules of solvent</p>                                                             | <p>As a hæmostatic in hæmorrhage due to superficial injuries and after tonsillectomy; epistaxis; external bleeding of hæmophilics; and in nasal, oral and dental surgery</p>                                                                     |



## HOW TO USE

## PACKINGS

One or more, taken with a little water, twice or thrice daily, after food

**'Tabloid' Reduced Iron**, bottles of 100

Half to one measureful (0.2 gm. to 0.4 gm.), daily, or more frequently, as indicated. A special measuring spoon is issued with each tube

**'Ryzamin-B,'** collapsible tubes of 15 grammes (1/2 oz. approx.) and 100 grammes, each with special nozzle

One, three times daily

**'Tabloid' Sodium Nitrite Compound**, bottles of 25 and 100

Orally, 0.5 mgm. to 5 mgm., three times a day. By intramuscular injection, 1 mgm. to 5 mgm., three times a day

**'Tabloid' Stilbœstrol** 0.5 mgm., 1 mgm. and 5 mgm., each strength in bottles of 25 and 100. **'Hypoloid' Stilbœstrol**, 1 mgm. in 1 c.c. and 5 mgm. in 1 c.c., each strength in boxes of 6 ampoules

The Sodium Tetraiodophenolphthalein and Acid Mixture, dissolved and mixed, provide a freshly precipitated suspension for oral administration

**'Stipolac,'** cartons containing one tube of each, and boxes containing 25 pairs

May be applied to the bleeding-point on gauze or on a pledget; by dropping on the part from a fine needle, or by means of an atomiser spray

**'Stypven,'** containers of 1 c.c. (0.1 mgm. Venom and Solvent) and 5 c.c. (0.5 mgm. Venom and Solvent)

| PRODUCT                                                                                                                                                                                                        | INDICATIONS                                                                                                                                                                                                                                                                                                                                                                                                                                |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>Sulphonamide-P</b><br/>(Sulphanilamide)</p> <p>'Tabloid' Sulphonamide-P, 0.5 gm. (for oral administration); 'Wellcome' Sulphonamide-P (for parenteral or oral administration)</p>                        | <p>Puerperal fever, complications of scarlet fever, erysipelas, tonsillitis, Ludwig's angina and other hæmolytic streptococcal infections; meningococcal, streptococcal and pneumococcal meningitis; gonococcal infections; <i>B. coli</i> and <i>Pr. ammoniæ</i> infections of the genito-urinary tract. May be of value in gas-gangrene, typhoid and paratyphoid fevers, <i>Br. abortus</i> infections and staphylococcal septicæmia</p> |
| <p>TRADE MARK 'Tannafax' BRAND</p> <p><b>Tannic Acid Jelly</b></p> <p>Tannic acid, with 0.5 per cent. phenol, in a water-soluble base</p>                                                                      | <p>Burns and scalds of every type (including electrical and chemical burns); superficial wounds and abrasions; bed-sores</p>                                                                                                                                                                                                                                                                                                               |
| <p>TRADE MARK 'Theoba' MARK</p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>R Theobrominæ, gr. 5<br/>Phenobarbitoni, gr. 1/2</p>                                                                                      | <p>Hypersensitive heart disease; angina pectoris and other conditions associated with high blood-pressure; intermittent claudication</p>                                                                                                                                                                                                                                                                                                   |
| <p><b>Three Bromides Effervescent</b></p> <p>TRADE MARK 'Tabloid' BRAND</p> <p>R Potassii Bromidi, 0.4 gramme<br/>Sodii Bromidi, 0.4 gramme<br/>Ammonii Bromidi, 0.2 gramme<br/>Salis Effervescentis, g.s.</p> | <p>As a sedative in conditions of nervous excitability; epilepsy; insomnia; and in sea-sickness and vomiting from reflex causes</p>                                                                                                                                                                                                                                                                                                        |
| <p><b>Vitamin B<sub>1</sub> Hydrochloride</b></p> <p>The pure crystalline anti-neuritic principle, with a potency of 333 International Units per mgm.</p>                                                      | <p>Beri-beri; polyneuritis of alcoholic, diabetic or post-infectious origin; and other conditions in which high Vitamin B<sub>1</sub> dosage is indicated</p>                                                                                                                                                                                                                                                                              |

## HOW TO USE

## PACKINGS

Orally, up to six or eight grammes daily. Parenterally, 0.5 gramme in 40 c.c. of normal saline. Intrathecal injection is also sometimes employed. (*For details of administration, precautions and contra-indications, see special literature*)

**'Tabloid' Sulphonamide-P,** bottles of 25 and 100. **'Wellcome' Sulphonamide-P,** bottles of 25 grammes and 100 grammes

Apply lightly, allow to dry, and bandage lightly or not at all. Do not apply oil or grease. In abraded wounds, etc., it is important to cleanse thoroughly the affected area before application

**'Tannafax,'** collapsible tubes of 20 gm. (3/4 oz. approx.) and 4 oz. (113 gm. approx.)

One or two, taken with a little water or milk

**'Tabloid' 'Theoba,'** bottles of 25 and 100

One to two, dissolved in half-a-tumblerful of water, after food

**'Tabloid' Three Bromides, Effervescent,** tubes of 25

Orally, 1 mgm. to 3 mgm., three times a day. By injection, 2 mgm. to 10 mgm.

**'Tabloid' Vitamin B<sub>1</sub> Hydrochloride,** 1 mgm., bottles of 25. **'Hypoloid' Vitamin B<sub>1</sub> Hydrochloride,** 2 mgm. in 1 c.c. and 10 mgm. in 1 c.c., each strength in boxes of 6 ampoules

## INDEX

PAGE

|                                                                                                           |    |
|-----------------------------------------------------------------------------------------------------------|----|
| Acetarsol, 'KHAROPHEN' ... ..                                                                             | 18 |
| Acetylsalicylic Acid ( <i>see</i> 'TABLOID,' 'EMPIRIN' COMPOUND,<br>page 10; 'TABLOID' 'EMOCIN,' page 10) |    |
| Adrenaline preparations ... ..                                                                            | 2  |
| 'ALEPOL' ... ..                                                                                           | 2  |
| ALKALINE COMPOUND, EFFERVESCENT, 'TABLOID' ... ..                                                         | 2  |
| Amœbicides ( <i>see</i> Carbarsone, page 8; 'KHAROPHEN,' page 18;<br>'QUINOXYL,' page 20)                 |    |
| AMYL NITRITE, 'VAPOROLE' ... ..                                                                           | 2  |
| Antimony ( <i>see</i> 'NEOSTAM,' page 20)                                                                 |    |
| Arsphenamines ( <i>see</i> 'NEOKHARSIVAN,' page 18;<br>'KHARSULPHAN,' page 18)                            |    |
| ASCORBIC ACID, 'TABLOID' ... ..                                                                           | 4  |
| 'AVENYL' ... ..                                                                                           | 4  |
| BARBITONE, 'TABLOID' ... ..                                                                               | 4  |
| ( <i>see also</i> 'TABLOID' PHENOBARBITONE, page 20)                                                      |    |
| 'BICREOL' ... ..                                                                                          | 4  |
| Bismuth preparations for injection ... ..                                                                 | 4  |
| BLAUD PILL, 'TABLOID' ... ..                                                                              | 6  |
| Bromides ( <i>see</i> 'TABLOID' THREE BROMIDES,<br>EFFERVESCENT, page 24)                                 |    |
| Burn dressing ( <i>see</i> 'TANNAFAX,' page 24)                                                           |    |
| Calciferol ... ..                                                                                         | 6  |
| ( <i>see also</i> 'TABLOID' CALCIUM GLUCONATE WITH<br>CALCIFEROL, page 6)                                 |    |
| Calcium Gluconate preparations ... ..                                                                     | 6  |
| Calcium Iodo-ricinoleate, 'IODICIN' ... ..                                                                | 16 |
| CALCIUM LÆVULATE, 'HYPOLOID' ... ..                                                                       | 6  |
| CALCIUM MANDELATE COMPOUND, 'WELLCOME' ... ..                                                             | 8  |
| Carbarsone ... ..                                                                                         | 8  |
| CAROTENE, 'TABLOID' ... ..                                                                                | 8  |
| Cevitamic Acid ( <i>see</i> 'TABLOID' ASCORBIC ACID, page 4)                                              |    |
| Chaulmoogric Esters ( <i>see</i> 'MOOGROL,' page 18; IODISED<br>'MOOGROL,' page 18)                       |    |
| Chiniofon, 'QUINOXYL' ... ..                                                                              | 20 |
| Codeine ( <i>see</i> 'TABLOID' 'EMPIRIN' COMPOUND WITH<br>CODEINE, page 10)                               |    |
| COD LIVER OIL WITH MALT EXTRACT, 'KEPLER' ... ..                                                          | 18 |
| Diethylstilbœstrol ( <i>see</i> Stilbœstrol, page 22)                                                     |    |
| 'DIGINUTIN' ... ..                                                                                        | 8  |
| DIGITALIS LEAF, 'TABLOID' ... ..                                                                          | 8  |
| DIGOXIN (B. W. & Co.) ... ..                                                                              | 10 |
| Diuretics ( <i>see</i> 'HYPOLOID' MERSALYL, page 18; 'TABLOID'<br>'THEOBA,' page 24)                      |    |
| 'EMOCIN,' 'TABLOID' ... ..                                                                                | 10 |
| 'EMPIRIN' COMPOUND, 'TABLOID' ... ..                                                                      | 10 |
| Ephedrine preparations ... ..                                                                             | 12 |



|                                                                                                                                                                                                                        | PAGE |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Epinephrine ( <i>see</i> Adrenaline preparations, <i>page</i> 2)                                                                                                                                                       |      |
| Ergometrine ... ..                                                                                                                                                                                                     | 12   |
| Ergotoxine Ethanesulphonate ... ..                                                                                                                                                                                     | 14   |
| 'ERNUTIN' ... ..                                                                                                                                                                                                       | 14   |
| ERYTHRITYL TETRANITRATE, 'TABLOID' ... ..                                                                                                                                                                              | 14   |
| 'ETHIDOL' ... ..                                                                                                                                                                                                       | 14   |
| 'EULYKOL' ... ..                                                                                                                                                                                                       | 14   |
| FERROUS CHLORIDE, CITRATED, 'TABLOID' ... ..                                                                                                                                                                           | 14   |
| GLYCERYL TRINITRATE, 'TABLOID' ... ..                                                                                                                                                                                  | 16   |
| Hæmostatics ( <i>see</i> 'STYFVEN,' <i>page</i> 22; 'WELLCOME' SOLUTION OF ADRENALINE TARTRATE, <i>page</i> 2)                                                                                                         |      |
| Hydnocarpus Oil ( <i>see</i> 'ALEPOL,' <i>page</i> 2; 'EULYKOL,' <i>page</i> 14)                                                                                                                                       |      |
| Hypnotics ( <i>see</i> 'TABLOID' BARBITONE, <i>page</i> 4; 'TABLOID' PHENOBARBITONE, <i>page</i> 20)                                                                                                                   |      |
| 'INFUNDIN' ... ..                                                                                                                                                                                                      | 16   |
| INSULIN, 'WELLCOME' ... ..                                                                                                                                                                                             | 16   |
| 'IODICIN' ... ..                                                                                                                                                                                                       | 16   |
| Iodine preparations ( <i>see</i> 'ETHIDOL,' <i>page</i> 14; 'IODICIN,' <i>page</i> 16)                                                                                                                                 |      |
| IODISED 'MOOGROL' ... ..                                                                                                                                                                                               | 18   |
| Iodophthalein ( <i>see</i> 'STIPOLAC,' <i>page</i> 22)                                                                                                                                                                 |      |
| Iron preparations ( <i>see</i> 'TABLOID' BLAUD PILL, <i>page</i> 6; 'TABLOID' FERROUS CHLORIDE, CITRATED, <i>page</i> 14; 'TABLOID' REDUCED IRON, <i>page</i> 22; 'TABLOID' CALCIUM GLUCONATE AND IRON, <i>page</i> 6) |      |
| 'KEPLER' COD LIVER OIL WITH MALT EXTRACT ... ..                                                                                                                                                                        | 18   |
| 'KHAROPHEN' ... ..                                                                                                                                                                                                     | 18   |
| 'KHARSULPHAN' ... ..                                                                                                                                                                                                   | 18   |
| MALT EXTRACT, 'KEPLER' COD LIVER OIL WITH ... ..                                                                                                                                                                       | 18   |
| Mandelic Acid ( <i>see</i> 'WELLCOME' CALCIUM MANDELATE COMPOUND, <i>page</i> 8)                                                                                                                                       |      |
| MERSALYL, 'HYPOLOID' ... ..                                                                                                                                                                                            | 18   |
| 'MOOGROL' ... ..                                                                                                                                                                                                       | 18   |
| 'NEOKHARSIVAN' ... ..                                                                                                                                                                                                  | 18   |
| 'NEOSTAM' ... ..                                                                                                                                                                                                       | 20   |
| 'NICAMIDE' ... ..                                                                                                                                                                                                      | 20   |
| Nicotinic Acid ... ..                                                                                                                                                                                                  | 20   |
| Nicotinic Acid Diethylamide, 'NICAMIDE' ... ..                                                                                                                                                                         | 20   |
| Nitroglycerin ( <i>see</i> 'TABLOID' GLYCERYL TRINITRATE, <i>page</i> 16)                                                                                                                                              |      |
| Œstrogen, synthetic ( <i>see</i> Stilbæstrol, <i>page</i> 22)                                                                                                                                                          |      |
| Oxytocics ( <i>see</i> Ergometrine, <i>page</i> 12; Ergotoxine Ethanesulphonate, <i>page</i> 14; 'ERNUTIN,' <i>page</i> 14; 'INFUNDIN,' <i>page</i> 16)                                                                |      |
| PHENOBARBITONE, 'TABLOID' ... ..                                                                                                                                                                                       | 20   |
| Phenylethyl hydnocarpate ( <i>see</i> 'EULYKOL,' <i>page</i> 14)                                                                                                                                                       |      |
| Pituitary (Posterior Lobe) Extract, 'INFUNDIN' ... ..                                                                                                                                                                  | 16   |
| PROTAMINE INSULIN, 'WELLCOME' ... ..                                                                                                                                                                                   | 16   |

|                                                                                                                                                                                             | PAGE |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Pyridine $\beta$ -Carboxylic Acid Diethylamide ( <i>see</i> 'NICAMIDE,' page 20)                                                                                                            |      |
| 'QUINOXYL' ... ..                                                                                                                                                                           | 20   |
| REDUCED IRON, 'TABLOID' ... ..                                                                                                                                                              | 22   |
| Russell Viper Venom, 'STYPVEN' ... ..                                                                                                                                                       | 22   |
| 'RYZAMIN-B' ... ..                                                                                                                                                                          | 22   |
| Sodium Hydnocarpate ( <i>see</i> 'ALEPOL,' page 2)                                                                                                                                          |      |
| SODIUM NITRITE COMPOUND, 'TABLOID' ... ..                                                                                                                                                   | 22   |
| Spirillicides ( <i>see</i> Bismuth preparations, page 4; 'NEOKHARSIVAN,' page 18; 'KHARSULPHAN,' page 18; 'AVENYL,' page 4)                                                                 |      |
| Stibamine Glucoside, 'NEOSTAM' ... ..                                                                                                                                                       | 20   |
| Stilboestrol ... ..                                                                                                                                                                         | 22   |
| 'STIPOLAC' ... ..                                                                                                                                                                           | 22   |
| 'STYPVEN' ... ..                                                                                                                                                                            | 22   |
| Sulphanilamide ( <i>see</i> Sulphonamide-P, page 24)                                                                                                                                        |      |
| Sulpharsphenamine, 'KHARSULPHAN' ... ..                                                                                                                                                     | 18   |
| Sulphonamide-P ... ..                                                                                                                                                                       | 24   |
| 'TANNAFAX' ... ..                                                                                                                                                                           | 24   |
| Tetraiodophenolphthalein, 'STIPOLAC' ... ..                                                                                                                                                 | 22   |
| Tetranitrin ( <i>see</i> 'TABLOID' ERYTHRITYL TETRANITRATE, page 14)                                                                                                                        |      |
| 'THEOBA,' 'TABLOID' ... ..                                                                                                                                                                  | 24   |
| Thiamin ( <i>see</i> Vitamin B <sub>1</sub> Hydrochloride, page 24)                                                                                                                         |      |
| THREE BROMIDES, EFFERVESCENT, 'TABLOID' ... ..                                                                                                                                              | 24   |
| Trinitrin ( <i>see</i> 'TABLOID' GLYCERYL TRINITRATE, page 16)                                                                                                                              |      |
| Urinary antiseptics ( <i>see</i> 'WELLCOME' CALCIUM MANDELATE COMPOUND, page 8; Sulphonamide-P, page 24)                                                                                    |      |
| Vaso-dilators ( <i>see</i> 'VAPOROLE' AMYL NITRITE, page 2; 'TABLOID' ERYTHRITYL TETRANITRATE, page 14; 'TABLOID' GLYCERYL TRINITRATE, page 16; 'TABLOID' SODIUM NITRITE COMPOUND, page 22) |      |
| Vitamin A ( <i>see</i> 'TABLOID' CAROTENE, page 8; 'KEPLER' COD LIVER OIL WITH MALT EXTRACT, page 18)                                                                                       |      |
| Vitamin B <sub>1</sub> Hydrochloride ... ..                                                                                                                                                 | 24   |
| ( <i>see also</i> 'RYZAMIN-B,' page 22)                                                                                                                                                     |      |
| Vitamin B <sub>2</sub> ( <i>see</i> Nicotinic Acid, page 20)                                                                                                                                |      |
| Vitamin C ( <i>see</i> 'TABLOID' ASCORBIC ACID, page 4)                                                                                                                                     |      |
| Vitamin D ( <i>see</i> Calciferol, page 6; 'TABLOID' CALCIUM GLUCONATE WITH CALCIFEROL, page 6; 'KEPLER' COD LIVER OIL WITH MALT EXTRACT, page 18)                                          |      |




**BURROUGHS WELLCOME & CO.**  
(THE WELLCOME FOUNDATION LTD.)  
**LONDON**

*Associated Houses:*

|          |          |          |              |       |
|----------|----------|----------|--------------|-------|
| NEW YORK | MONTREAL | SYDNEY   | CAPE TOWN    | MILAN |
|          | BOMBAY   | SHANGHAI | BUENOS AIRES |       |



Reduced facsimiles



Hæmolytic  
Streptococci  
in Pus

# SULPHONAMIDE-P

## (SULPHANILAMIDE)

TRADE MARK 'TABLOID' BRAND SULPHONAMIDE-P

*For Oral administration*

TRADE MARK 'WELLCOME' BRAND SULPHONAMIDE-P

*For Parenteral or Oral administration*

*(See pages 24 and 25)*

HARLEY STREET,  
LONDON. W.1.

R  
'Tabloid' 'Empirin' Comp.  
c Codeina  
(Original package)  
Sig. Sum. i ter die  
post cibos

W.G.